






In this Section:


- Stripping
- Stock piling/Importing
- Spreading (Topsoiling)
- Testing
- Amending
 - Fertilizers
 - Compost
































We know more about the movement of celestial bodies than about the soil underfoot.


Leonardo Da Vinci








Stripping Depth! % Clay 20-30 10-20 o-30 cm 0-10 cm cm cm Ea 26.8 21 21 22.9 Lb 38 17.5 21.6 25.7 Nd 18.5 35.3 42.5 32.1 Ng 18.5 35.3 42.5 32.1 Sb 11.5 28 11.5 17 Tc 19.5 32.5 39 39











Maintaining Soil Life in Topsoil Stock Piles

To maintain native soil life (microbial, fungal, mycorrhizae and other life) we need:

- Grow plants on the pile
- Best would be: native plants (native annuals???)


Minimum
Standards 2 and 1



MS-2 (Topsoil stockpiles)

- Soil stockpiles and borrow areas shall be stabilized or protected with sediment trapping measures
- Applies to on and off-site stockpiles and borrow areas





Stockpiling Topsoil: ESC Standards and Specifications 3.30



Stabilized and protect in accordance with MS-2



Side slopes shall not exceed 2:1



Perimeter controls should be placed immediately



 Seeding within 7 days if left dormant for more than 14 days in accordance with MS-1



Seeding to Maintain Life in Topsoil Stock Piles

Seeding is generally done with native annuals (Std. & Specs. 3.31)

- Roots provide oxygen, carbon (organic matter) and other nutrients to microorganisms in the soil
- In the past literature suggested a cover crop of

Buckwheat

buckwheat

Remember

Seeding is not just done to meet some regulation or to prevent erosion!

Plants helps in keeping the soil (biologically but also physically) healthy and improve the chances of revegetation success once the topsoil is reapplied to the site.






Module 3c

Spreading (Topsoiling)











Do not apply when soil is wet or frozen



 Minimum 2" deep on 3:1 or steeper slopes 4" on flatter slopes



Avoid irregular surface (ponding)



 May need some compaction for good bonding with subsoil (avoid over-compaction)

(Top)Soil and the BMP Clearinghouse · Sheet flow Surface Grass Channels Vegetated Roof Infiltration • Dry swale • Extended Detention Pond (2) Subsurface · Permeable Pavement Infiltration Infiltration Bioretention No · Rainwater Harvesting Infiltration • Filtering Practices Wet Swale Ponding Constructed Wetland • Wet Pond • Extended Detention Pond (1)



2. Sheet Flow

- Conserved Open Space → native soil = No disturbance!
- Vegetated Filter Strip → All soil types, no fill, yes topsoil

	Topsoil Requirements							
	Soil type	Loamy sand/sandy loam						
	Clay content	<5%						
When needing to import topsoil	pН	6-7						
	Organic matter	≥ 2 ⁰ ⁄ ₀						
	Salt	< 500 ppm						
	Depth	6-8" lightly compacted						



3. Grass Channels

- Can be used on all soil types, with or without compost amendments
- Hydrologic soil groups C and D require compost amendments to improve performance
- No topsoil requirements in Standards and Specifications, but no reason not to use topsoil
- Protection of soil (infiltration ability) stressed!





10. Dry Swale

- Can be used on all soil types
- Hydrologic soil groups C and D require an underdrain
- Dry swales require a 4" **topsoil** layer on top of an engineered soil medium (Table 10.4) of at least 18" that is placed on top of choker stone layers on native soil
- Alternatively, use: 18" of sand on top of which is placed a 4" combination (mix?) of topsoil and leaf compost covered by turf





10. Dry Swale

Table 10.4

Topsoil Requirements						
Soil type	Loamy sand/sandy loam					
Clay content	<5%					
pН	6-7					
Organic matter	≥ 2%					
Salt	< 500 ppm					
Depth	6-8" lightly compacted					





15. Extended Detention Pond Level 2

- No recommendations for the use of topsoil
- Recommend using ESC Standards and Specifications for the use of topsoil (buffers, etc)









SW BMPs Conclusion


- For most BMPs soil at siting is more important →
 infiltration (Module 7)
- Topsoil is utilized in two ways:
 - 1. Non-disturbed/restored
 - 2. To cover up an engineered medium

















- Soil texture, pH and nutrient content should be determined before application of lime and fertilizer
- Soil samples are sent for testing
- Small fee
- Saves time, money
- Will help to establish vegetation correctly





So, what's in that soil test report?

- Available from various labs. Usually cost \$10 to \$20 per sample for simple tests; can be much expensive.
 Be alert for "voodoo prescriptions"!
- Typically report pH, OM, and dilute acid extractable P, Ca, Mg, K and Na. You can also ask for soluble salts and extractable micronutrients (Zn, Mn, Fe, etc.)





A Typical Report

SAMPLE HISTORY										
Sample Field ID ID		LAST CROP	LAST LIME APPLICATION		SOIL INFORMATION					
		Name	Yield	Months Free.	Tons/Acre	SMU-1 %	SMU-2 %	SMU-3 %	Yield Estimate	Productivity Group
OCF11	4463	Orchaedgraan/Fescue-Clover Pasture (40)		18+		40B2 100				III
	LAB TEST RESULTS (see Note 1)									

ELECTRICAL (ACTIONS)											
Analysis	P (lb/A)	K (lb/A)	Ca (lb/A)	Mg (lb/A)	Zn (p	ppm) Ma	(ppm)	Си (ррг	n) Fe (ppu	a) B (ppm)	S.Salts (ppm)
Result	9	95	1409	209	1.	.2 1	0.3	0.3	4.4	0.5	
Rating	L+	M-	M+	H+	su	FF S	UPP	SUPP	SUFI	SUFF	
Analysis	Soil pH	Buffer Index	EstCE (meq/100			Base Sat. (%)	C	a Sat. (%)	Mg Sat. (%)	K Sat. (%)	Organic Matter (%)
Result	E 7	6 21	г.	c 20	-	70.0		60 E	15.2	2.2	2 6

FERTILIZER AND LIMESTONE RECOMMENDATIONS

Lbs/acre = lb/A ppm (pounds per million) \rightarrow 1 acre soil = 2 million lbs Zn 1.2 ppm \rightarrow 2.4 lb/A





Example of Fertilizer Recommendation

FERTILIZER AND LIMESTONE RECOMMENDATIONS

Crop: Orchardgrass/Fescue-Clover Pasture (40)

Lime, T	ONS/AC	Fertilizer, lb/A						
Amount	Type	N	P205	K20				
1	AG	50	40	50				

825. If stand contains less than 25 per cent clover, apply 40-60 lbs N/A.

131. If additional production is needed later on, apply 40 to 60 lbs/A of N during the grazing season. If you are planning to overseed a legume into the stand, omit the N recommendation.

122. P2O5 and K2O recommendations are for annual application. However, rates can be doubled and applied every other year if desired.





Soil Amendments




Need to be based on soil analysis!







Soil Organic Matter (Compost)



Also known as *Humus*. It is the dark brown to black complex decomposition product of organic matter turnover in soils. It is typically reported as *organic matter* content in soil testing programs. In a soil profile it typically occurs in the O profile (O = Organic) and is leached out into the A profile.



Soil Organic Matter (Compost)

Organic Matter in the soil:

Opens soil up: increases infiltration



• Binds soil particles together: decreases erosion



Binds nutrients: improves fertility



Feeds microorganisms: improves overall soil health





